



TE

L:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBQ004M101 100µg
Histone Cluster 1, H4a (HIST1H4A)
Organism Species: Multi-species
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Ser2~Gly103 (Accession # P62806), with N-terminal His-Tag.

Host: *E. coli*

Subcellular Location: Nucleus. Chromosome.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).


Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% sucrose, 0.01% sarcosyl.

Predicted isoelectric point: 11.2

Predicted Molecular Mass: 12.9kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)





TE L:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Note: Full length bovine HIST1H4A is identical in sequence to mouse HIST1H4A and rat HIST1H4A. 100% cross-reactivity of HIST1H4A was observed among bovine, mouse and rat.

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.



TE L:4006-871-227 Web:www.ybio.net
Email:shybio@126.com

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MG HH HHH HS GSE F- SG RG KG GK G LG KG GA KRH R KVLRD NIQ GI TKPAIRR LAR
RGGVKRISGL IYEETRGVLK VFLENVIRDA VTYTEHAKRK TVTAMDVVYA LKRQGRTLYG
FGG