



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB504Mu01 10 μ g

Recombinant Tumor Necrosis Factor Receptor Superfamily,

Member 1B (TNFRSF1B)

Organism Species: *Mus musculus* (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: His303~Ala474

Tags: N-terminal His-Tag

Accession: P25119

Host: *E. coli*


Subcellular Location: Membrane; Single-pass type I membrane protein.

Purity: >95%

Endotoxin Level: <1.0EU per 1 μ g (determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 5.6



The possible reasons that the actual band size differs from the predicted are as follows:

Predicted Molecular Mass: 19.4kDa

Accurate Molecular Mass: 32kDa as determined by SDS-PAGE reducing conditions. Applications: SDS-PAGE; WB; ELISA; IP.



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

(May be suitable for use in other assays to be determined by the end user.)

Note:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.



TEL:4006-871-227 Web:www.ybio.net

Email:shybio@126.com

4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.

5. Polymerization of the target protein: Dimerization, multimerization etc

[USAGE]

Reconstitute in sterile ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

```
HVPDEKSQL DAVGLEQQHL LTTAPSSSSS SLESSASAGD RRAPPGGHPQ ARVMAEAQGF  
QEARASSRIS DSSHGSHGTH VNVTCIVNVC SSSDHSSQCS SQASATVGDV DAKPSASPKD  
EQVPPSQEEC PSQSPCETTE TLQSHEKPLP LGVPDMGMKP SQAGWFDQIA VKVA
```

[REFERENCES]

1. Lewis M., *et al.* (1991) Proc. Natl. Acad. Sci. U.S.A. 88:2830-2834.
2. Goodwin R.G., *et al.* (1991) Mol. Cell. Biol. 11:3020-3026.
3. Hurle B., *et al.* (1998) Genomics 52:79-89.



TEL:4006-871-227 Web:www.ybio.net

Email:shybio@126.com

4. Van Ess P.J., *et al.* (2002) J. Pharmacol. Exp. Ther. 300:824-830.