



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB830Hu01 10 μ g

Recombinant Transglutaminase 2, Tissue (TGM2)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Met1~Ala687 (Accession # P21980),
with N-terminal His-Tag.

Host: *E. coli*

Purity: >95%

Endotoxin Level: <1.0EU per 1 μ g
(determined by the LAL method).

Formulation: Supplied as lyophilized form in
PBS, pH7.4, containing 1mM DTT, 5% trehalose,
0.1% sarcosyl and preservative.

Predicted isoelectric point: 5.3

Predicted Molecular Mass: 78.8kDa

The possible reasons that the actual band size differs from the predicted are as follows:

Accurate Molecular Mass: 100kDa as determined by SDS-PAGE reducing
conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.



TEL:4006-871-227 Web:www.ybio.net

Email:shybio@126.com

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MGHHHHHSGSEF-MAEELVLERC DLELETNGRD HHTADLCREK LVVRRGQPFW
LTLHFEGRNY EASVDSLTFV VVTGPAPSQE AGTKARFPLR DAVEEGDWTA TVVDQQDCTL
SLQLTPANA PIGLYRLSLE ASTGYQGSSF VLGHFILLFN AWC PADAVYL DSEERQEVV
LTQQGFIYQG SAKFIKNIPWNFGQFEDGIL DICLILLDVN PKFLKNAGRD CSRRSSPVVY
GRV V S G M V N C ND D Q G V LL G R WD N N Y G D G V S PM S W I G S V D I LR RW K N H G C Q
RVKYGQCWVF AAVACTVLRC LGIPTRVVTN YNSAHDQNSN LLIEYFRNEF GEIQGDKSEM
IWNFHCVVES WM TRPDLQPGYEGWQALDPTPQEKSEGTCCGPVPVRAIK
EGDLSTKYDA PFVFAEVNAD VVDWIQQDDG SVHKSINRSL IVGLKISTKS VGRDEREDIT
HTYKYPEGSS EEREAFTAN HLNKLAKEE TGMAMRIRVG QSMNMGSDFD VFAHITNNTA
EYVCRLLC ARTVSYNGIL GPCGTYKLL NLNLEPFSEK SVPLCILYK YRDCLTESNL
IKVRALLVEP VINSYLLAER DLYLENPEIK IRILGEPKQK RKLVAEVS LQ NPLPVALEGC
TFTVEGAGLT EEQKTVEIPD PVEAGEEVKV RMDLLPLHMG LHKLVVNFES DKLKAVKGFR
NVIIGPA

[REFERENCES]



TEL:4006-871-227 Web:www.ybio.net

Email:shybio@126.com

1. Gentile V., *et al.* (1991) J. Biol. Chem. 266:478-483.
2. Fraij B.M., *et al.* (1992) J. Biol. Chem. 267:22616-22623.
3. Van Damme P., *et al.* (2012) Proc. Natl. Acad. Sci. U.S.A. 109:12449-12454.
4. Pinkas D.M., *et al.* (2007) PLoS Biol. 5:E327-E327.