

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA139Ca01 100µg

Recombinant Tumor Necrosis Factor Related Apoptosis Inducing Ligand (TRAIL)

Organism Species: Canis familiaris; Canine (Dog)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Thr36~Gly281

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: F1PWB1

Host: E. coli

Purity: >95%

Endotoxin Level: <1.0EU per 1 µ g (determined by the

LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4

containing 5% trehalose, 0.01% sarcosyl.


Predicted isoelectric point: 8.6

Predicted Molecular Mass: 32.6kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]





Web:www.ybio.net Email:shybio@126.com

Reconstitute in sterile PBS, pH7.2-pH7.4.



Email:shybio@126.com

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

TRELK QMQDKYSQSG IACFLKEDDI PWDPSDEESM NNPCWQVKWQ LRQFVRKMIL

KTYEETIPTA PEKQLNIPYV VSDRGSQRVA AHITGTSRRS MFPIPSSKND KALGHKINSW

DSTRKGHSFL NNLHLRNGEL VIHQRGFYYI YSQTYFRFQE PEEIPTGQNR KRNKQMVQYI

YKHTSYPDPI LLMKSARNSC WSKDSEYGLY SIYQGGIFEL KENDRIFVSV SNEQLIDMDQ

EASFFGAFLI G