



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB959Hu01 50µg
Recombinant Cluster Of Differentiation 34 (CD34)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Ser32~Thr290

Tags: N-terminal His-Tag

Accession: P28906

Host: *E. coli*

**Subcellular Location: Membrane; Single-pass
type I membrane protein.**

Purity: >95%


**Endotoxin Level: <1.0EU per 1µg (determined by
the LAL method).**

**Formulation: Supplied as lyophilized form in PBS
pH7.4, containing 1mM DTT, 5% trehalose, 0.05%
sarcosyl and preservative.**

Predicted isoelectric point: 6.1

Predicted Molecular Mass: 29.0kDa

Accurate Molecular Mass: 35kDa as determined by SDS-PAGE reducing



The possible reasons that the actual band size differs from the predicted are as follows:

conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.**
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.**
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.**



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.

5. Polymerization of the target protein: Dimerization, multimerization etc.



TEL:4006-871-227 Web:www.ybio.net

Email:shybio@126.com

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

SLDNNGTAT PELPTQGTF S NVSTNVS YQE TTPSTL GST SLHPV SQHGN EATTNITETT
VKFTSTSVIT SVYGNTN SSV QSQTSVISTV FTTPANVSTP E TTLKPSLSP GNVSDLSTTS
TSLATSPTKP YTSSSPILSD IKAEIKCSGI REVKLTQ GIC LEQNK TSSCA EFKKDRGEGL
ARVLCGEEQA DADAGA QVCS LLLAQSEVRP QCLLLVLANR TEISSKLQLM KKHQSDLKKL
GILDFTEQDV ASHQSYSQKT

[REFERENCES]

1. Simmons D.L., *et al.* (1992) J. Immunol. 148:267-271.
2. Satterthwaite A.B., *et al.* (1992) Genomics 12:788-794.
3. Nakamura Y., *et al.* (1993) Exp. Hematol. 21:236-242.
4. Sutherland D.R., *et al.* (1988) Leukemia 2:793-803.