

YBD744Mu01 50µg

Recombinant Lipolysis Stimulated Lipoprotein Receptor (LSR) Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES



[<u>PROPERTIES</u>]

Residues: Ser35~Glu205

Tags: Two N-terminal Tags, His-tag and GST-tag Accession: 099KG5

Host: E. coli

Subcellular Location: Cell membrane; Single-pass

type I membrane protein.

Purity: >95%

Endotoxin Level: <1.0EU per $1\,\mu\,g$ (determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris, 500mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 4.7

Predicted Molecular Mass: 48.7kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

[<u>USAGE</u>]

Reconstitute in ddH₂O.



[<u>STORAGE AND STABILITY</u>]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCES</u>]

The sequence of the target protein is listed below. SAIQVT VPDPYHVVIL FQPVTLHCTY QMSNTLTAPI VIWKYKSFCR DRVADAFSPA SVDNQLNAQL AAGNPGYNPY VECQDSVRTV RVVATKQGNA VTLGDYYQGR RITITGNADL TFEQTAWGDS GVYYCSVVSA QDLDGNNEAY AELIVLGRTS EAPELLPGFR AGPLE

[<u>REFERENCES</u>]

Lin Q., et al. (1996) Submitted to the EMBL/GenBank/DDBJ databases.
Yen F.T., et al. (2008) J. Biol. Chem. 283:25650-25659.
Stenger C., et al. (2012) J. Neurochem. 123:467-476.
Sohet F., et al. (2015) J. Cell Biol. 208:703-711.