



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBC539Ra01 100µg

Recombinant Intestinal Cell Kinase (ICK)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Met1~Phe284

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: Q62726

Host: *E. coli*

Subcellular Location: Cytoplasm.

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).


Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose, 0.01% sarcosyl.

Predicted isoelectric point: 9.3

Predicted Molecular Mass: 36.9kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)



[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

**MNRYTTIKQL GDGTYGSVLL GRSIESGELI AIKKMKRKFY SWEECMNLRE VKSLKKNLHA
NIVKLKEVIR ENDHLYFIFE YMKENLYQLI KERNKLPES AIRNIMYQIL QGLAFIHKHG
FFHRDLKPEN LLCMGPELVK IADFGlareI RSRPPYTDYV STRWYRAPEV LLRSTNYSSP
IDVWAVGCIM AEVYTLRPLF PGASEIDTIF KICQVLGTPK KTDWPEGYQL SSAMNFIWPQ
CIPNNLKTli PNASSEAIQL LRDLLQWDPK KRPTASQALR YPYF**