

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBE909Hu01 100µg

Recombinant Histone Deacetylase 9 (HDAC9)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Pro23~Ala343

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: Q9UKVO

Host: E. coli

Subcellular Location: Nucleus.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the

LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4,

containing 5% trehalose, 0.01% sarcosyl.


Predicted isoelectric point: 9.6

Predicted Molecular Mass: 40.1kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]





Web:www.ybio.net Email:shybio@126.com

Reconstitute in sterile PBS, pH7.2-pH7.4.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below. PLDLRTDL RMMMPVVDPV VREKQLQQEL LLIQQQQQIQ KQLLIAEFQK QHENLTRQHQ AQLQEHIKEL LAIKQQQELL EKEQKLEQQR QEQEVERHRR EQQLPPLRGK DRGRERAVAS TEVKQKLQEF LLSKSATKDT PTNGKNHSVS RHPKLWYTAA HHTSLDQSSP PLSGTSPSYK YTLPGAQDAK DDFPLRKTAS EPNLKVRSRL KQKVAERRSS PLLRRKDGNV VTSFKKRMFE VTESSVSSSS PGSGPSSPNN GPTGSVTENE TSVLPPTPHA EQMVSQQRIL IHEDSMNLLS LYTSPSLPNI TLGLPAVPSQ LNA