

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBC128Mu01 10µg

Recombinant Gremlin 1 (GREM1)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

11th Edition (Revised in May, 2016)

## [PROPERTIES]

Source: Prokaryotic expression.

Host: E. coli

Residues: Lys25~Asp184 Tags: N-terminal His-Tag

Tissue Specificity: Lung, Kidney. Subcellular Location: Secreted.

**Purity: >92%** 

**Traits:** Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 1mM DTT, 5% trehalose, 0.01%

sarcosyl and Proclin300.

Original Concentration: 200ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; Reporter Assays; Purification;

Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.1

Predicted Molecular Mass: 23.9kDa

Accurate Molecular Mass: 27kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:



TEL:4006-871-227 Web:www.ybio.net

Email:shybio@126.com

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.



TEL: 4006-871-227 Web: www. ybio. net

Email:shybio@126.com

- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

#### [USAGE]

Reconstitute in PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### [SEQUENCE]

# [ IDENTIFICATION ]



871-227 Web:www.ybio.net

Email:shybio@126.com

Figure 1. Gene Sequencing (Extract)



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.




Figure 2. SDS-PAGE