

YBA133Bo01 100µg Recombinant Tumor Necrosis Factor Alpha (TNFa) Organism Species: Bos taurus; Bovine (Cattle) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES


9th Edition (Revised in Jul, 2013)

[<u>PROPERTIES</u>]

Residues: Metl~Leu234 (Accession # Q06599), with two N-terminal Tags, His-tag and T7-tag. Host: *E. coli* Subcellular Location: Secreted. Membrane; Single-pass type I membrane protein. Purity: >95% Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% sucrose, 0.01% sarcosyl. Predicted isoelectric point: 6.3 Predicted Molecular Mass: 29.4kDa Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[<u>USAGE</u>]



TEL: 4006-871-227

Web:www.ybio.net Email:shybio@126.com Reconstitute in sterile PBS, pH7.2-pH7.4.



[<u>STORAGE AND STABILITY</u>]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37° C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCES</u>]

The target protein is fused with two N-terminal Tags, His-tag and T7-tag, its sequence is listed below. MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- MSTKSMIRDV ELAEEVLSEK AGGPQGSRSC LCLSLFSFLL VAGATTLFCL LHFGVIGPQR EEQSPGGPSI NSPLVQTLRS SSQASSNKPV AHVVADINSP GQLRWWDSYA NALMANGVKL EDNQLVVPAD GLYLIYSQVL FRGQGCPSTP LFLTHTISRI AVSYQTKVNI LSAIKSPCHR ETPEWAEAKP WYEPIYQGGV FQLEKGDRLS AEINLPDYLD YAESGQVYFG IIAL