

YB90061Rb01 Interleukin 15 (IL15)

Organism: Oryctolagus cuniculus (Rabbit) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC **PROCEDURES**

5th Edition (Revised in January, 2013)

Rabbit IL15

kDa 94 66.2

45

33

26

20

14.4

[DESCRIPTION]

Protein Names: Interleukin 15

Synonyms: IL15 Species: Rabbit

Size: 100)g

Source: Escherichia coli-derived Subcellular Location: Secreted.

[PROPERTIES]

Residues: Asn49~Pro162 (Accession # Q3Y5G8), with N-

terminal His-Tag.

Grade & Purity: >95%, 18kDa as determined by SDS-PAGE 15% SDS-PAGE

reducing conditions.

Formulation: Supplied as lyophilized form in PBS, pH 7.4,

containing 5% sucrose, 0.01% sarcosyl.

Endotoxin Level: <1.0 EU per 1µg (determined by the LAL

method).

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Predicted Molecular Mass: 14.6kDa

Predicted isoelectric point: 5.6

Email:shybio@126.com

[PREPARATION]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGSEF -NW HDVISDLKRI EDLIKSIHID ATLYTESDAH PNCKVTAMKC FLLELRVISH ESRNMDINET VQNLIILANT SLSSKGNVTE SGCKECEELE EKNITEFLQS FIHIVQMFIN SP