
TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA285M101 100µg Histone H3 (H3) Organism Species: Multi-species Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES



Predicted Molecular Mass: 16.9kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

9th Edition (Revised in Jul, 2013)



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com Note: Full length human histone H3 is identical in sequence to mouse histone 3 and rat histone H3. 100% cross-reactivity of Histone H3 was observed among human, mouse and rat.

[<u>USAGE</u>]

Reconstitute in sterile PBS, pH7.2-pH7.4.



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months. Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGSEF-MARTKQTARK STGGKAPRKQ LATKAARKSA PATGGVKKPH RYRPGTVALR EIRRYQKSTE LLIRKLPFQR LVREIAQDFK TDLRFQSSAV MALQEACEAY LVGLFEDTNL CAIHAKRVTI MPKDIQLARR IRGERA

[REFERENCES]

1. Zhang Q., et al. (2013) Cell Res. 23:225-241. 2. Krishnan S., Trievel R.C. (2013) Structure 21:98-108. 3. Cai L., et al. (2013) Mol. Cell 49:571-582. 4. Ballare C., et al. (2012) Nat. Struct. Mol. Biol. 19:1257-1265.