



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBG901Hu01 100µg

Recombinant Voltage Dependent Anion Channel Protein 1 (VDAC1)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Ala2~Ala283

Tags: Two N-terminal Tags, His-tag and GST-tag

Accession: P21796

Host: *E. coli*

Subcellular Location: Mitochondrion outer membrane.

Cell membrane.

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose, 0.01% sarcosyl.

Predicted isoelectric point: 8.6


Predicted Molecular Mass: 60.6kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

**AVPPTYADL GKSARDVFTK GYGFLIKLD LTKSENGLE FTSSGSANTE TTKVTGSLET
KYRWTEYGLT FTEKWNTDNT LGTEITVEDQ LARGLKLTFD SSFSPNTGKK NAKIKTGYKR
EHINLGCDMD FDIAGPSIRG ALVLGYEGWL AGYQMNFFETA KSRVTQSNFA VGYKTDEFQL
HTNVNDGTEF GGSYQKVNK KLETAVNLAW TAGNSNTRFG IAAKYQIDPD ACFSKVNNS
SLIGLGYTQT LKPGIKLTLA ALLDGKNVNA GGHKLGLGLE FQA**