

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBD066Hu01 100μg Recombinant Apelin (APLN) Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

# [PROPERTIES]

Residues: Ser24~Phe77 (Accession # Q9ULZ1),

with N-terminal His-Tag.

Host: E. coli

Subcellular Location: Secreted.

**Purity: >95%** 

Endotoxin Level: <1.0EU per 1µg (determined by the


LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris, 500mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and

preservative.

Predicted isoelectric point: 11.9 Predicted Molecular

Mass: 7.6kDa



15% SDS-PAGE

The possible reasons that the actual band size differs from the predicted are as follows:  $\frac{1}{2} \left( \frac{1}{2} \right) = \frac{1}{2} \left( \frac{1}{2} \right) \left($ 

Accurate Molecular Mass: 10kDa as determined by SDS-PAGE reducing conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.) Note:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

TEL:4006-871-227 Web:www.ybio.net

Email:shybio@126.com

# [USAGE]

Reconstitute in ddH2O.

# [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

# [SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MGHHHHHHSG SEF- SLMPLPD GNGLEDGNVR HLVQPRGSRN GPGPWQGGRR

KFRQRPRLS HKGPMPF

# [REFERENCES]

- 1. Tatemoto K., et al. (1998) Biochem. Biophys. Res. Commun. 251:471-476.
- 2. Habata Y., et al. (1999) Biochim. Biophys. Acta 1452:25-35.
- 3. Cayabyab M., et al. (2000) J. Virol. 74:11972-11976.
- 4. Foldes G., et al. (2003) Biochem. Biophys. Res. Commun. 308:480-485.