

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YB91497Hu01 100µq Protein Disulfide Isomerase A3 (PDIA3) Organism: Homo sapiens (Human) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

6th Edition (Revised in March, 2013)

kDa

26

20

14.4

15% SDS-PAGE

[PROPERTIES]

Residues: Asp162~Tyr416 (Accession # P30101),

with N-terminal His-Tag.

Host: E. coli

Subcellular Location: Endoplasmic reticulum

lumen. Melanosome.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS,

pH7.4, containing 5% sucrose, 0.01% sarcosyl.

Predicted isoelectric point: 5.8

Predicted Molecular Mass: 30.8kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MG HHHH HHSG SEF- DSFSE AHSE FLKAAS NLRD NYR FAHTNVE SLVNEY DDNG

EGIILFRPSH LTNKFEDKTV AYTEQKMTSG KIKKFIQENI FGICPHMTED NKDLIQGKDL

LIAYYDVDYE KNAKGSNYWR NRVMMVAKKF LDAGHKLNFA VASRKTFSHE LSDFGLESTA

GEIPVVAIRT AKGEKFVMQE EFSRDGKALE RFLQDYFDGN LKRYLKSEPI PESNDGPVKV

VVAENFDEIV NNENKDVLIE FYAPWCGHCK NLEPKY