



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

**YBC591Ra01  
100ug**

**Recombinant Ubiquitin Specific Peptidase 1 (USP1)**

**Organism Species: Rattus norvegicus (Rat)**

*Instruction  
manual*

**FOR IN VITRO USE AND RESEARCH USE ONLY**

**NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES**

10th Edition (Revised in Jan,  
2014)

## **[ PROPERTIES ]**

**Residues: Met1~His198**

**Tags: Two N-terminal Tags, His-tag and T7-tag**

**Accession: Q569C3**

**Host: *E. coli***

**Subcellular Location: Nucleus.**

**Purity: >90%**

**Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).**


**Formulation: Supplied as lyophilized form in 20mM**

**Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM**

**DTT, 0.01% sarcosyl, 5% trehalose, and preservative.**

**Predicted isoelectric point: 6.7**

**Predicted Molecular Mass: 26.0kDa**



The possible reasons that the actual band size differs from the predicted are as follows:

**Accurate Molecular Mass: 31kDa as determined by SDS-PAGE reducing conditions.**

**Applications: SDS-PAGE; WB; ELISA; IP.**

**(May be suitable for use in other assays to be determined by the end user.)**

**Note:**



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

1. **Splice variants:** Alternative splicing may create different sized proteins from the same gene.
2. **Relative charge:** The composition of amino acids may affects the charge of the protein.
3. **Post-translational modification:** Phosphorylation, glycosylation, methylation etc.
4. **Post-translation cleavage:** Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. **Polymerization of the target protein:** Dimerization, multimerization etc.

## **[ USAGE ]**

Reconstitute in sterile ddH<sub>2</sub>O.

## **[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCES ]**

The sequence of the target protein is listed below.

MPGVIPSESN GLSRGSPSKK NRLSLKFFQK KETKRALDFT DSQEDEEKAS EYRGSEIDQV  
VPAAQSSPVS CEKRENLLPF VGLNNLGNTC YLNSILQVLY FCPGFKAGVK HLFNIISRKK  
EALKDDSIQK DKGSKCEDPL ASYELICSLQ SLIISVEQLQ ASFLLNPEKY TDELATQPRR  
LLNLTRELNP MYEGYLQH