



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB750Ca01 100 μ g

Tumor Necrosis Factor Ligand Superfamily, Member 13 (TNFSF13)

Organism Species: *Canis familiaris*; Canine (Dog)

*Instruction
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Glu92~Leu250 (Accession # B5B3U5), with two
N-terminal Tags, His-tag and S-tag.

Host: *E. coli*

Purity: >95%

Endotoxin Level: <1.0EU per 1 μ g
(determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4,
containing 5% sucrose, 0.01% sarcosyl.

Predicted isoelectric point: 8.9


Predicted Molecular Mass: 23.8kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.





[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and S-tag, its sequence is listed below.

```
MHHHHHSSG LVPRGSGMKE TAAAKFERQH MDSPDLGTDD DDKAMADIGS EF-  
EAWENGERS RRKRAALHK QKKKHSVLHL VPINITSKED SDVTEVMWQP ALKRGRGLEA  
Q G Y V V R V W D S G I Y L L Y S Q V L F H D V T F T M G Q V V S R E G Q G R Q E T L F R C I R S M  
PSNPDWAYNS CYSAGVFHLH QGDILSVTIP RGRAKLSLSP HGTFLGFVKL
```