

YB97833Hu01 Iduronate-2-Sulfatase (IDS)

Organism: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC **PROCEDURES**

5th Edition (Revised in January, 2013)

[DESCRIPTION]

Protein Names: Iduronate-2-Sulfatase

Synonyms: IDS, SIDS

Species: Human

Size: 100,g

Source: Escherichia coli-derived Subcellular Location: Lysosome.

[PROPERTIES]

Residues: Arg95~Pro289 (Accession # P22304), with N-

terminal His-Tag.

Grade & Purity: >95%, 25kDa as determined by SDS-PAGE

reducing conditions.

Formulation: Supplied as lyophilized form in PBS, pH 7.4,

containing 5% sucrose, 0.01% sarcosyl.

Endotoxin Level: <1.0 EU per 1µg (determined by the LAL

method).


Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Predicted Molecular Mass: 23.8kDa

Predicted isoelectric point: 6.1

Human IDS





[PREPARATION]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGSEF-RRPDTT RLYDFNSYWR VHAGNFSTIP QYFKENGYVT MSVGKVFHPG ISSNHTDDSP YSWSFPPYHP SSEKYENTKT CRGPDGELHA NLLCPVDVLD VPEGTLPDKQ STEQAIQLLE KMKTSASPFF LAVGYHKPHI PFRYPKEFQK LYPLENITLA PDPEVPDGLP PVAYNPWMDI RQREDVQALN ISVPYGPIP