TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB028Bo01 100µg

Native Albumin (ALB)

Organism Species: Bos taurus; Bovine (Cattle)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Host: Native

Source: Bovine Plasma

Subcellular Location: Secreted.

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in 50mM TRIS, 200mM

NaCl Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[RELEVANCE]

The albumins are a family of globular proteins, the most common of which is serum albumin. The albumin family consists of all proteins that are watersoluble, are moderately soluble in concentrated salt solutions, and experience heat denaturation. Albumins are commonly found in blood plasma, and are unique from other blood proteins in that they are not glycosylated. Substances containing albumins, such as egg white, are called albuminoids. Its main function is to regulate the colloidal osmotic pressure of blood.



[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[REFERENCES]

- 1. Bujacz A. (2012) Acta Crystallogr. D Biol. Crystallogr. 68:1278-1289.
- 2. Majorek K.A., et al. (2012) Mol. Immunol. 52:174-182.
- 3. Hilger C., et al. (2001) Clin. Exp. Immunol. 123:387-394.
- 4. Hirayama K., et al. (1990) Biochem. Biophys. Res. Commun. 173:639-646.