YB99111Hu01
DNA Repair Protein RAD50 (RAD50)
Organism: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

[DESCRIPTION]

Protein Names: DNA Repair Protein RAD50

Synonyms: RAD50

Species: Human

Size: 100μg

Source: *Escherichia coli*-derived

Subcellular Location: Nucleus. Chrom
telomere.

[PROPERTIES]

Residues: Val400~Ala595 (Accession # Q92878)
with N-terminal His-Tag.

Grade & Purity: >95%, 27kDa as determined by
SDS-PAGE reducing conditions.

Formulation: Supplied as lyophilized form in PBS, pH
7.4, containing 5% sucrose, 0.01% sarcosyl.

Endotoxin Level: <1.0 EU per 1μg (determined by
the LAL method).

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Predicted Molecular Mass: 24.6kDa

Predicted isoelectric point: 6.6

[PREPARATION]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8^{O}C for one month.

Aliquot and store at -80^{O}C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37^{O}C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MGHHHHHHSGSEF- V RERQEGEAKT ANQLMNDFAE KETLKQKQID EIRDKKTGLG

RIIELKSEIL SKKQNELKNV KYELQQLEGS   SDRILELDQE LIKAERELSK AEKNSNVETL

KMEVISLQNE KADLDRTLRK LDQEMEQLNH HTTTRTQMEM LTKDKADKDE QIRKIKSRHS

DELTSLLGYF PNKKQLEDWL HSKSKEINQT RDRLA