TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBC565Hu01 100µg

Recombinant Eukaryotic translation initiation factor 2D (EIF2D)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

## [ PROPERTIES ]

Residues: Met1~Leu117

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: P41214

Host: E. coli

Subcellular Location: Cytoplasm.

Purity: >95%

Endotoxin Level: <1.0EU per 1 µ g (determined by the LAL

method).

Formulation: Supplied as lyophilized form in PBS, pH7.4,

containing 5% trehalose, 0.01% sarcosyl.


Predicted isoelectric point: 7.9

Predicted Molecular Mass:

16.6kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)



## [ USAGE ]

Reconstitute in sterile PBS, pH7.2-pH7.4.

## [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## [ SEQUENCES ]

The sequence of the target protein is listed below. MFAKAFRVKS NTAIKGSDRR KLRADVTTAF PTLGTDQVSE LVPGKEELNI VKLYAHKGDA VTVYVSGGNP ILFELEKNLY PTVYTLWSYP DLLPTFTTWP LVLEKLVGGA DLMLPGL