

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA394Rb01 100µg

Recombinant Tissue Factor Pathway Inhibitor (TFPI) Organism Species: Oryctolagus cuniculus (Rabbit)

Instruction manual

kD:

66.

45

33

26

20

14.

15% SDS-PAGE

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Lys39~Gly275 (Accession # P19761), with two N-

Host: E. coli

Subcellular Location: Secreted.

terminal Tags, His-tag and GST-tag.

Purity: >95%

Endotoxin Level: <1.0EU per 1 µ g

(determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS,

pH7.4, containing 1mM DTT, 5% trehalose,

sarcosyl and preservative.

Predicted isoelectric point: 7.9

Predicted Molecular Mass: 59.2kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]



Web:www.ybio.net Email:shybio@126.com

Reconstitute in sterile PBS, pH7.2-pH7.4.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and GSTtag, its sequence is listed below.

MSPILGYWKI KGLVQPTRLL LEYLEEKYEE HLYERDEGDK WRNKKFELGL EFPNLPYYID

GDVKLTQSMA IIRYIADKHN MLGGCPKERA EISMLEGAVL DIRYGVSRIA YSKDFETLKV

DFLSKLPEML KMFEDRLCHK TYLNGDHVTH PDFMLYDALD VVLYMDPMCL DAFPKLVCFK

KRIEAIPQID KYLKSSKYIA WPLQGWQATF GGGDHPPKSD GSTSGSGHHH HHHSAGLVPR

GS TA I G M K E T AA A K F E R Q H M DS P D L G T L E V LF Q G P L G S E F - K P PL Q K P T H S F C

AMKVDDGPCR AYIKRFFFNI LTHQCEEFIY GGCEGNENRF ESLEECKEKC ARDYPKMTTK


LTFQKGKPDF CFLEEDPGIC RGYITRYFYN NQSKQCERFK YGGCLGNLNN FESLEECKNT

CENPTSDFQV DDHRTQLNTV NNTLINQPTK APRRWAFHGP SWCLPPADRG LCQANEIRFF

YNAIIGKCRP FKYSGCGGNE NNFTSKKACI TACKKGFIPK SIKGG

[REFERENCES]

- 1. Wesselschmidt R.L., et al. (1990) Nucleic Acids Res. 18:6440-6440.
- 2. Warn-Cramer B. J., et al. (1992) Nucleic Acids Res. 20:3548-3548.



- 3. Belaaouaj A., *et al.* (1993) Thromb. Res. 69:547-553.
- 4. Luo J.G., et al. (2009) Zhonghua Xin Xue Guan Bing Za Zhi 37:1113-1118.