



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

## NADH 脱氢酶 5(ND5)重组蛋白

适用生物 Rattus norvegicus (Rat , 大鼠)  
产品应用 SDS-PAGE; WB; ELISA; IP. 规格 50ug  
分子量 16.8kDa 价格 请咨询当地代理商  
纯度 > 95%  
来源 原核表达

## Recombinant NADH Dehydrogenase 5 (ND5)

FOR IN VITRO AND RESEARCH USE ONLY, NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES!

Organism species	Rattus norvegicus (Rat)
Product No.	YBS952Ra01
Source	Prokaryotic expression
Host	E. coli
Purity	> 95%
UOM	50ug
Predicted Molecular Mass	16.8kDa
Predicted isoelectric point	n/a
Applications	SDS-PAGE; WB; ELISA; IP.
Endotoxin Level	<1.0EU per 1μg (determined by the LAL method)

Subcellular Location	n/a
Residues	Arg425~Lys562 (Accession # AP_004902.1) with N-terminal His-Tag
Formulation	Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose, 0.01% sarcosyl

## SEQUENCES



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MGHHHHHSGSEF-RIIFYVTMTK PRYSPLITIN ENNPNLINPI KRLALGSILA GFLISLNIPP TNIQILTI  
PH LKITALLITI LGFAIALELN NLTLNLSISK PTKLSSFSTS LGYYPPIIPR IIPQKTLNSS YKLSLNLLD  
L ISEKTIPK

## USAGE

Reconstitute in sterile PBS, pH7.2-pH7.4.

## STORAGE AND STABILITY

**Storage:** Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## About the MARKER (complimentary)

**Effective Size Range:** 10kDa to 70kDa.

**Protein bands:** 10kDa, 14kDa, 18kDa, 22kDa, 26kDa, 33kDa, 44kDa and 70kDa.

**Double intensity bands:** The 26kDa, 18kDa, 10kDa bands are at double intensity to make location and size approximation of proteins of interest quick and easy.

**Ready-to-use:** No need to heat, dilute or add reducing agents before use.