



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBG476Mu01 100 μ g

Recombinant Histidine Decarboxylase (HDC)

Organism Species: *Mus musculus* (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Tyr341~Ser597

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: P23738

Host: *E. coli*

Purity: >95%

Endotoxin Level: <1.0EU per 1 μ g
(determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS,
pH7.4, containing 5% trehalose, 0.01% sarcosyl.

Predicted isoelectric point: 9.7

The possible reasons that the actual band size differs from the predicted are as follows:

Predicted Molecular Mass: 32.5kDa

Accurate Molecular Mass: 38kDa as determined by SDS-PAGE reducing
conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note:



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.



TEL:4006-871-227 Web:www.ybio.net

Email:shybio@126.com

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

YLRHANS GAA TDFMHWQIPL SRRFRS IKLW FVIRSF GVKV N LQAHVRHGTE MAKYFESLVR
SDPSFEIPAK RHLGLVVFRL KGPNCLETESV LKEIAKAGQL FLIPATIQDK LIIRFTVTSQ
FTTKEDILRD WHLIQEAAANL VLSQHCTSQP SPRAKNVIPP PPGTRGLSLE SVSEGGDDPA
QARKIIKQPG ASLARREGGS DLETMPDPFD DCFSEEAPNT TKHKLSSFLF SYLSVQNRRK
TTRSLSCNSV PMSAQKS