TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB947Ra01 100µg

Recombinant Acrosin (ACR)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

## [PROPERTIES]

Residues: Ile43~Gly291

Tags: N-terminal His-Tag

Accession: P29293

Host: E. coli **Purity: >95%** 

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris,

500mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01%

sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 8.0


Predicted Molecular Mass: 29.1kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

# [USAGE]

Reconstitute in ddH2O.





## [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

#### [SEQUENCES]

The sequence of the target protein is listed below.

IVGGQTSS PGAWPWMVSL QIFTSHNSRR YHACGGSLLN SHWVLTAAHC FDNKKKVYDW RLVFGAHEIE YGRNKPVKEP QQERYVQKIV IHEKYNAVTE GNDIALLKVT PPVTCGDFVG PGCLPHFKSG PPRIPHTCYV TGWGYIKDNA PRPSPVLMEA RVDLIDLDLC NSTQWYNGRV TSTNVCAGYP EGKIDTCQGD SGGPLMCRDS VDSPFVIVGI TSWGVGCARA KRPGVYTATW DYLDWIASKI G

#### [ REFERENCES ]

- 1. Klemm U., et al. (1991) Biochim. Biophys. Acta 1090:270-272.
- 2. Kremling H., et al. (1991) DNA Seq. 2:57-60.
- 3. Kwitek A.E., et al. (2004) Genome Res. 14:750-757.
- 4. Szpirer C., et al. (1998) Mamm. Genome 9:721-734.