YBF541Mu01 100µg Recombinant Leukocyte Cell Derived Chemotaxin 2 (LECT2) Organism Species: Mus musculus (Mouse) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

kDa

70 44

33

26

22

18

14

10

15% SDS-PAGE

[PROPERTIES]

Residues: Met1~Leu151

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: 088803

Host: E. coli

Subcellular Location: Secreted.

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM

Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 9.4

Predicted Molecular Mass: 20.2kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile ddH₂O.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

MIPTTILISA ALLSSALAGP WANICASKSS NEIRTCDSYG CGQYSAQRTQ RHHPGVDVLC SDGSVVYAPF TGKIVGQEKP YRNKNAINDG IRLSGRGFCV KIFYIKPIKY KGSIKKGEKL GTLLPLQKIY PGIQSHVHVE NCDSSDPTAY L

[REFERENCES]

- 1. Yamagoe S., et al. (1998) Gene 216:171-178.
- 2. Shukunami C., et al. (1999) J. Biochem. 125:436-442.
- 3. Okumura A., et al. (2009) Biosci. Trends 3:139-143.
- 4. Ovejero C., et al. (2004) Hepatology 40:167-176.