L:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB039Ra01 100µg Recombinant Surfactant Associated Protein D (SPD) Organism Species: Rattus norvegicus (Rat) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

70 Residues: Ala20~Phe374 (Accession # P35248), with N-44 terminal His-Taq. Host: E. coli 33 Subcellular Location: Secreted, extracellular space, 26 extracellular matrix. surface film. 22 **Purity: >95%** 18 Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). 14 Formulation: Supplied as lyophilized form in PBS, pH7.4, 10 containing 5% sucrose, 0.05% sarcosyl. Predicted isoelectric point: 7.1 15% SDS-PAGE Predicted Molecular Mass: 37.0kDa Accurate Molecular Mass: 14&44kDa as determined by SDS-PAGE reducing conditions.

Applications: SDS-PAGE; WB; ELISA; IP.


(May be suitable for use in other assays to be determined by the end user.)

The possible reasons that the actual band size differs from the predicted are as follows: Note:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.







9th Edition (Revised in Jul, 2013)

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGSEF-A EMKTLSORSI TNTCTLVLCS PTENGLPGRD GRDGREGPRG EKGDPGLPGPMGLSGLPGPRGPVGPKGENGSAGEPGPKGERGLVGPPGSP G I S G PA G K E G PS G K Q G N I G P QG K P G P K G E A GP K G E V G A P G MQ G S A G A K G P AG P K G E R G A P GE Q G A P G N A G AA G PA G PA G P QG A P G S R G P P GL K G D R G A P G DRGIKGESGL PDSAALRQQM EALNGKLQRL EAAFSRYKKA ALFPDGQSVG DKIFRAANSE EPFEDAKEMC RQAGGQLASP RSATENAAVQ QLVTAHSKAA FLSMTDVGTE GKFTYPTGEA LVYSNWAPGE PNNNGGAENC VEIFTNGQWN DKACGEQRLV ICEF

[REFERENCES]

- 1. King B.A., Kingma P.S. (2011) Am. J. Respir. Cell Mol. Biol. 44:709-715.
- 2. Yamashita H., et al. (2010) Radiat Oncol 5:32-32.
- 3. Foreman M.G., et al. (2011) Am. J. Respir. Cell Mol. Biol. 44:316-322.
- 4. Stapleton R.D., et al. (2010) Chest 138:568-577.