



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA481Mu01 50µg

Recombinant Lymphotoxin Beta Receptor (LTbR)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

11th Edition (Revised in May, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Pro128~Tyr395

Tags: Two N-terminal Tags, His-tag and GST-tag

Tissue Specificity: Lung, Liver, Kidney, Pancreas.

Subcellular Location: Membrane; Single-pass type I membrane protein.

Purity: >92%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; Purification; Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.4

Predicted Molecular Mass: 58.4kDa

Accurate Molecular Mass: 47kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
PGM SCVYLDNECV HCEEERLVLC
QPGTEAEVTD EIMDTDVNCV PCKPGHFQNT SSPRARCQPH TRCEIQGLVE
AAPGTSYSDT ICKNPPEPGA MLLLAILLSL VLFLLFTTVL ACAWMRHPSL
CRKLGTLTKR HPEGEESPPC PAPRADPHFP DLAEP LLPMS GDLSPSPAGP
PTAPSLEEVS LQQQSPLVQA RELEAEPGEH GQVAHGANGI HVTGGSVTVT
GNIYIYNGPV LGGTRGPGDP PAPPEPPYPT PEEGAPGPSE LSTPY
```



TEL:4006-

871-

227 Web:www.ybio.net E

mail:shybio@126.com

[IDENTIFICATION]




Figure 1. SDS-PAGE