TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBG438Mu01 10µg Recombinant Calbindin(CALB) **Organism Species: Mus musculus (Mouse)** Instruction manual

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

kDa

#### [PROPERTIES]

| Residues: Glu3~Asn261 (Accession # P12658), with two N-      |              | 94<br>66.2 |
|--|--------------|------------|
| terminal Tags, His-tag and GST-tag.                          | -            | 45         |
| Host: <i>E. coli</i>   | -            | 33         |
| Purity: >95%   | -            | 26         |
| Endotoxin Level: <1.0EU per 1 $\mu$ g (determined by the LAL | 20           | 00         |
| method).   |              | 20         |
| Formulation: Supplied as lyophilized form in PBS, pH7.4,     | 14.          | 14.4       |
| containing 5% sucrose, 0.01% sarcosyl.                       |              |            |
| Predicted isoelectric point: 5.3                             | 15% SDS-PAGE |            |
| Predicted Molecular Mass: 61.5kDa                            |              |            |
| Applications: SDS-PAGE; WB; ELISA; IP.                       |              |            |

(May be suitable for use in other assays to be determined by the end user.)

## [USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.



#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

# [<u>SEQUENCES</u>]

The target protein is fused with two N-terminal Tags, His-tag and GST-tag, its sequence is listed below.

MSPILGYWKI KGLVQPTRLL LEYLEEKYEE HLYERDEGDK WRNKKFELGL EFPNLPYYID GDVKLTQSMA IIRYIADKHN MLGGCPKERA EISMLEGAVL DIRYGVSRIA YSKDFETLKV DFLSKLPEML KMFEDRLCHK TYLNGDHVTH PDFMLYDALD VVLYMDPMCL DAFPKLVCFK KRIEAIPQID KYLKSSKYIA WPLQGWQATF GGGDHPPKSD GSTSGSGHHH HHHSAGLVPR GSTAIGMKET AAAKFERQHM DSPDLGTLEV LFQ GPLGS- ESHLQSSL ITASQFFEIW LHFDADGSGY LEGKELQNLI QELLQARKKA GLELSPEMKS FVDQYGQRDD GKIGIVELAH VLPTEENFLL LFRCQQLKSC EEFMKTWRKY DTDHSGFIET EELKNFLKDL LEKANKTVDD TKLAEYTDLM LKLFDSNNDG KLELTEMARL LPVQENFLLK FQGIKMCGKE FNKAFELYDQ DGNGYIDENE LDALLKDLCE KNKQELDINN ITTYKKNIMA LSDGGKLYRT DLALILSAGD N

## [REFERENCES]

- 1. Godin J.D., et al. (2012) Dev. Cell 23:729-744.
- 2. Chen W.V., et al. (2012) Neuron 75:402-409.
- 3. Edqvist P.H., et al. (2012) PLoS ONE 7:e42488-e42488.
- 4. Peng C., et al. (2012) J. Biol. Chem. 287:32861-32873.