



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YB91370Ra01

Lactate Dehydrogenase A (LDHA)


Organism: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES

3th Edition (Revised in February, 2012)



[DESCRIPTION]

Protein Names: Lactate Dehydrogenase A

Gene Names: LDHA, Ldh-1, Ldh1

Size: 100 μ g

Source: Recombinant

Expression Host: *E.coli*

Subcellular Location: Cytoplasm.

[PROPERTIES]

Residues: Met1~Phe332 (Accession # P04642), with a N-terminal His-tag.

Grade & Purity: >97%, 38 kDa as determined by SDS-PAGE reducing conditions.

Form & Buffer: Supplied as lyophilized form in PBS, pH 7.4.

Endotoxin Level: <1.0 EU per 1 μ g (determined by the LAL method).

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Predicted Molecular Mass: 38 kDa

15% Tris-glycine SDS-PAGE



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

[**PREPARATION**]

Reconstitute in PBS.

[**STORAGE AND STABILITY**]

Storage: Store at 4°C for short time storage (1-2 weeks). Aliquot and store at -20°C or -80°C for long term storage. Avoid repeated freeze/thaw cycles.

Valid period: 12 months stored at -80°C.

[**BACKGROUND**]

The target protein is fused with a His-tag and its sequence is listed below. The first Met is an initiator amino acid. Moreover, Gly and Ser are added to improve the flexibility of N-terminus at both ends of the His-tag, which will increase the chelating ability of the tag to Ni-Sepharose during purification.

MGHHHHHSGSEF-MAALKDQLIV NLLKEEQVPQ NKITVVG VGA VGMACAISIL MKDLADELAL VDVIEDKLKG
EMMDLQHGSL FLKTPKIVSS KDYSVTANSK LVIITAGARQ QEGESRLNLV QRNVNIFKFI IPNVVKYSPQ
CKLLIVSNPV DILTYVAWKI SGFPKNRVIG SGCNLD SARF RYLMGERLGV HPLSCHGWVL GEHGDSSVPV
WSGVNVAGVS LKSLNPQLGT DADKEQWKDV HKQVVD SAYE VIKLKG YTSW AIGLSVADLA ESIMKNLRRV
HPISTMIKGL YGIKEDVFLS VPCILGQNGI SDVVKVTLTP DEEARLKKSA DTLWGIQKEL QF