

YBC628HuO1 10µg Recombinant Microseminoprotein Beta (MSMb) Organism Species: Homo sapiens (Human) Instruction manual

2013)

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul,

[<u>PROPERTIES</u>]

Residues: Asn19~Ile114 (Accession # P08118), with two N-terminal Tags, His-tag and S-tag. Host: *E. coli* Subcellular Location: Secreted. Purity: >95% Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 1mM DTT, 5% trehalose, 0.01% sarcosyl and preservative. Predicted isoelectric point:

The possible reasons that the actual band size differs from the predicted are as follows: 5.6 Predicted Molecular Mass: 16.7kDa



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Accurate Molecular Mass: 22kDa as determined by SDS-PAGE reducing conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com The target protein is fused with two N-terminal Tags, His-tag and S-tag, its sequence is listed below. MHHHHHHSSG LVPRGSGMKE TAAAKFERQH MDSPDLGTDD DDKAMADIGS EF- NA SCYFIPNEGV PGDSTRKCMD LKGNKHPINS EWQTDNCETC TCYETEISCC TLVSTPVGYD KDNCQRIFKK EDCKYIVVEK KDPKKTCSVS EWII

[REFERENCES]

1. Mbikay M., et al. (1987) DNA 6:23-29.

2. Ulvsbaeck M., et al. (1989) Biochem. Biophys. Res. Commun. 164:1310-1315.

3. Green C.B., et al. (1990) Biochem. Biophys. Res. Commun. 167:1184-1190.

4. Nolet S., et al. (1991) Biochim. Biophys. Acta 1089:247-249.