



YBA788Ra01

100µg Recombinant Programmed Cell Death Protein 1 Ligand 1
(PDCD1LG1)

Organism Species: *Rattus norvegicus* (Rat)

*Instruction
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

11th Edition (Revised in May,
2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Ser34~Val241

Tags: Two N-terminal Tags, His-tag and GST-tag

Purity: >95%

Traits: Freeze-dried powder

Buffer formulation: 100mM NaHCO₃, 500mM NaCl, pH8.3,
containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl,
5%Trehalose and Proclin300.

Original Concentration: 200ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; ReporterAssays;

Purification; Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)




Figure 2. SDS-PAGE



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Predicted isoelectric point:

8.2 Predicted Molecular Mass:

53.7kDa

Accurate Molecular Mass: 64kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in 100mM NaHCO₃, 500mM NaCl (pH8.3) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

SNVIMEC RFPVEQRLLD
 LALVYWEKE DKEVIQFVEG EEDLKPQHSS FRGRAFLPKD QLLKGNVAVLQ
 ITDVKLQDAG VYCCMISYGG ADYKRITLKV NAPYRKINQR ISMDPATSEH
 ELMCQAEQYP EAEVIWTNSD HQSLSGETTV TTSQTEEKLL NVTSLRVNA
 TANDVFHCTF WRVHSGENHT AELIPELPV PRLPHNRTHW V

[IDENTIFICATION]




Figure 1. Gene Sequencing (Extract)




Figure 2. SDS-PAGE



TEL:4006-871-

227 Web:www.ybio.net Email:shybio@126.com



EL:4006-871-
227 Web:www.ybio.
net Email:shybio
@126.com