TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB786Mu01 100ug

Recombinant Receptor Interacting Serine Threonine Kinase 2 (RIPK2)

Organism Species: Mus musculus (Mouse)

Instruction manua1

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

## [ PROPERTIES ]

Residues: Met1~Leu294

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: P58801

Host: E. coli

Subcellular Location: Cytoplasm.

Purity: >95%

Endotoxin Level: <1.0EU per 1 µ g (determined by the

LAL method).

Formulation: Supplied as lyophilized form in 20mM

Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTI


0.01% sarcosyl, 5% trehalose, and preservative.


Predicted isoelectric point: 6.3

Predicted Molecular Mass:

37. 4kDa

Applications: SDS-PAGE; WB; ELISA; IP.





(May be suitable for use in other assays to be determined by the end user.)

## [ USAGE ]

Reconstitute in sterile ddH2O.

## [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at  $37^{\circ}$ C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## [ SEQUENCES ]

The sequence of the target protein is listed below. MNGDAICSAL PPIPYHKLAD LHYLSRGASG TVSSARHADW RVRVAVKHLH IHTPLLDSER NDILREAEIL HKARFSYILP ILGICNEPEF LGIVTEYMPN GSLNELLHRK TEYPDIAWPL RFRILHEIAL GVNYLHNMNP PLLHHDLKTQ NILLDNEFHV KIADFGLSKW RMMSLSQSRS YKSAPEGGTI IYMPPENYEP GQKSRASVKH DIYSYAVIMW EVLSRKQPFE EVTNPLQIMY SVSQGHRPDT SEENLPFDIP HRGLMISLIQ SGWAQNPDER PSFLKCLIEL EPVL