



YBA548Mu01 50 μ g

Recombinant Intercellular Adhesion Molecule 1 (ICAM1)

Organism Species: *Mus musculus* (Mouse)

*Instruction
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan,
2014)

[PROPERTIES]

Residues: Val46~Leu301

Tags: N-terminal His-Tag

Accession: P13597


Host: *E. coli*

Subcellular Location: Membrane; Single-pass
type I membrane protein.

Purity: >95%

Endotoxin Level: <1.0EU per 1 μ g (determined by
the LAL method).

Formulation: Supplied as lyophilized form in
20mM Tris, 150mM NaCl, pH8.0, containing 1mM
EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and



The possible reasons that the actual band size differs from the predicted are as follows:
preservative.

Predicted isoelectric point:

4.8 Predicted Molecular Mass:

29.5kDa



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Accurate Molecular Mass: 36kDa as determined by SDS-PAGE reducing conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in sterile ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

VNCSS SCKEDLSLGL ETQWLKDELE SGPNWKLFEL SEIGEDSSPL CFENCQTVQS



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

SASATITVYS FPESVELRPL PAWQQVGKDL TLRCHVDGGA PRTQLSAVLL RGEEILSRQP
VGGHPKDPKE ITFTVLASRG DHGANFSCRT ELDLRPQGLA LFSNVSEARS LRTFDLPATI
PKLDTPDLLE VGTQQLFCS LEGLFPASEA RIYLELGGQM PTQESTNSSD SVSATALVEV
TEEFDRTLPL RCVLELADQI L

[REFERENCES]

1. Horley K. J., *et al.* (1989) EMBO J. 8:2889-2896.
2. Siu G., *et al.* (1989) J. Immunol. 143:3813-3820.
3. Ballantyne C. M., *et al.* (1989) Nucleic Acids Res. 17:5853-5853.
4. Ballantyne C. M., *et al.* (1992) Genomics 14:1076-1080.