



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA099Rb01 10 μ g

Recombinant Matrix Metalloproteinase 13 (MMP13)

Organism Species: *Oryctolagus cuniculus* (Rabbit)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Glu70~Thr245

Tags: N-terminal His-Tag

Accession: O62806

Host: *E. coli*

**Subcellular Location: Secreted, extracellular space,
extracellular matrix.**

Purity: >95%

**Endotoxin Level: <1.0EU per 1 μ g (determined by the LAL
method).**

**Formulation: Supplied as lyophilized form in 10mM
PBS, pH7.4, containing 1mM DTT, 5% trehalose,
0.01% sarcosyl and preservative.**

Predicted isoelectric point: 5.9


Predicted Molecular Mass: 20.9kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile ddH₂O.





[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

**E MQSFFGLEVT GKDDNTLAI MKQPRCGVPD VGEYNVFPRT LKWSQTNLTY RIVNYTPDLT
HSEVEKAFKK AFKVWSDVTP LNFTRIHNGT ADIMISFGTK EHGDFYPPFDG PSGLLAHAFP
PGPNYGGDAH FDDDETWTSS SKGYNLFLVA AHEFGHSLGL DSKDPGALM FPIYT**

[REFERENCES]

- 1. Vincenti M.P., *et al.* (1998) *Biochem. J.* 331:341-346.**
- 2. Nagase H., Woessner JF (1999) *J. Biol. Chem.* 274 (31): 21491 - 4.**
- 3. Maruyama K., Sugano S (1994) *Gene* 138 (1-2): 171 - 4.**
- 4. Gomis-Rüth FX., *et al.* (1997) *J. Mol. Biol.* 264 (3): 556 - 66.**