

YBA051Ra01 50µg

Recombinant Insulin Like Growth Factor 2 (IGF2)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

kDa

70

33 26

22

18

14

10

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Ala25~Glu91 (Accession # P01346), with Nterminal His-Tag.

Host: E. coli

Subcellular Location: Secreted.

Purity: >95%

Endotoxin Level: <1.0EU per 1 µ g

(determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4,

containing 1mM DTT, 5% trehalose, 0.05% sarcosyl and 15% SDS-PAGE

preservative.

Predicted isoelectric point: 6.6

Predicted Molecular Mass: 9.0kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]



Reconstitute in sterile PBS, pH7. 2-pH7. 4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSG SEF- AYRPSE TLCGGELVDT LQFVCSDRGF YFSRPSSRAN RRSRGIVEEC CFRSCDLALL ETYCATPAKS E

[REFERENCES]

- 1. Cornish J., et al. (2007) Am. J. Physiol. 292:E117-E122.
- 2. Whitfield H. J., et al. (1984) Nature 312:277-280.
- 3. Frunzio R., et al. (1986) J. Biol. Chem. 261:17138-17149.
- 4. Soares M.B., et al. (1985) Nucleic Acids Res. 13:1119-1134.
- 5. Chiariotti L., et al. (1988) Mol. Endocrinol. 2:1115-1126.
- 6. Soares M.B., et al. (1986) J. Mol. Biol. 192:737-752.