

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YB81689Po02 100µg

Retinol Binding Protein 7, Cellular (RBP7)

Organism: Sus scrofa; Porcine (Pig)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

7th Edition (Revised in May, 2013)

[PROPERTIES]

Residues: Metl~Ala134 (Accession # B9P3U3), with two N-

terminal Tags, His-tag and T7-tag.

Host: E. coli

Purity: >95%

Endotoxin Level: <1.0EU per 1μg (determined by the LAL

method).

Formulation: Supplied as lyophilized form in PBS, pH7.4,

containing 5% sucrose, 0.01% sarcosyl.

Predicted isoelectric point: 7.8


Predicted Molecular Mass: 19.0kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile PBS, pH7. 2-pH7. 4.





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at $37^{\circ}\mathrm{C}$ for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and T7-tag, its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGS-MPADLSGTWN LVSSDNFEGY MLALGIDFAT RKIAKLLKPQ KVIEQNGDSF TIHTNSSLRN YLVKFKVGEE FDEDNKGLDN RKCKSLVTWD NDRLTCVQKG EKKNRGWTHW IEGDELHLEM FCEGQVCKQT FQRA