



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBE237Mu01 50μg

Hyaluronoglucosaminidase 2 (HYAL2)

Organism Species: *Mus musculus* (Mouse)

*Instruction
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul,
2013)

[PROPERTIES]

Residues: Leu23~Gly447 (Accession # 035632), with N-terminal His-Tag.

Host: *E. coli*

Subcellular Location: Cell membrane; Lipid-anchor,
GPI-anchor.

Purity: >95%

Endotoxin Level: <1.0EU per 1μg (determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4,
containing 5% sucrose, 0.01% sarcosyl.


Predicted isoelectric point:

The possible reasons that the actual band size differs from the predicted are as follows:

7.3 Predicted Molecular Mass:

50.2kDa

Accurate Molecular Mass: 44kDa as determined by SDS-PAGE reducing
conditions. Applications: SDS-PAGE; WB; ELISA; IP.





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

(May be suitable for use in other assays to be determined by the end user.)

Note:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MG H H H H H S G S E F - L K P T A P P I F T G R P F V V A W N V P T Q E C A P R H K V P L D L R A F
DVKATPNEG F FNQNIITFY DRLGLYPRFD AAGTSVHGGV PQNGSLCAHL PMLKESVERY
I Q T Q E P G G L A V I D W E E W R P V W V R N W Q E K D V Y R Q S S R Q L V A S R H P D W P S D R
V M K Q A Q Y E F E F A R Q F M L N T L R Y V K A V R P Q H L W G F Y L F P D C Y N H D Y V Q N W



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

ESYTGRCPDV EVARNDQLAW LWAESTALFP SVYLDETLAS SVHSRNFVSF RVREALRVAH
THHANHALPV YVFTRPITYTR GLTGLSQVDL ISTIGESAAL GSAGVIFWGD SEDASSMETC
QYLKNYLTQL LVPYIVNVSW ATQYCSWTQC HGHGRCVRRN PSANTFLHLN ASSFRLVPGH
TPSEPQLRPE GQLSEADLNY LQKHFRQCQY LGWGGEQCQR NYKGAAG

[REFERENCES]

1. Strobl B., *et al.* (1998) Genomics 53:214-219.
2. Rai S.K., *et al.* (2001) Proc. Natl. Acad. Sci. U.S.A. 98:4443-4448.
3. Chang N.-S. (2002) BMC Cell Biol. 3:8-8.
4. Duh F.M., *et al.* (2005) Retrovirology 2:59-59.