TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA551Mu01 50µg

Recombinant Fibroblast Growth Factor 2, Basic (FGF2)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES.

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Pro10~Ser154

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: P15655

Host: F coli

Subcellular Location: Secreted, Nucleus,

Purity: >95%

Endotoxin Level: <1.0EU per 1µg

(determined by the LAL method). **Formulation**: Supplied as

lyophilized form in

100mM NaHCO3, 500mM NaCl, pH8.3, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and

preservative.

Predicted isoelectric point: 9.7 Predicted Molecular Mass: 20.1kDa

Accurate Molecular Mass: 20&23kDa as determined by SDS-PAGE


reducing conditions.

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note: The possible reasons that the actual band size differs from the predicted are as follows:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.





[USAGE]

Reconstitute in sterile ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

P ALPEDGGAAF PPGHEKDPKR LYCKNGGEEL RIHPDGRVDG **VREKSDPHVK** LOLOAFERGY VSIKGYCANR YLAMKEDGRI. LASKCYTEEC EFFERLESNN YNTYRSRKYS SWYVALKRTG QYKLGSKTGP GQKAILFLPM SAKS

[REFERENCES]

- 1. Hebert J.M., et al. (1990) Dev. Biol. 138:454-463.
- 2. Kurtz A., et al. (1997) Oncogene 14:2671-2681.
- 3. Yamanaka Y., et al. (2011) Mol. Cell. Neurosci. 46:200-212.
- 4. Chikazu D., et al. (2001) J. Bone Miner. Res. 16:2074-2081.