

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBJ560Ra01 100µg

Recombinant Branched Chain Alpha-Ketoacid Dehydrogenase Kinase (BCKDK)

Organism Species: Rattus norvegicus (Rat)

> Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

> 10th Edition (Revised in Jan, 2014)

## [ PROPERTIES ]

Residues: Arg44~Thr223

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: Q00972

Host: E. coli

Subcellular Location: Mitochondrion.

Purity: >95%

Endotoxin Level: <1.0EU per 1μg (determined by

the LAL method).


Formulation: Supplied as lyophilized form in PBS,

pH7. 4, containing 5% trehalose, 0.01% sarcosyl.

Predicted isoelectric point:

9.0 Predicted Molecular Mass:

24. 5kDa



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

## [ USAGE ]

Reconstitute in sterile PBS, pH7.2-pH7.4.

## [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## [ SEQUENCES ]

The sequence of the target protein is listed below. RERSKTV TSFYNQSAID VVAEKPSVRL TPTMMLYSGR SQDGSHLLKS GRYLQQELPV RIAHRIKGFR SLPFIIGCNP TILHVHELYI RAFQKLTDFP PIKDQADEAQ YCQLVRQLLD DHKDVVTLLA EGLRESRKHI EDEKLVRYFL DKTLTSRLGI RMLATHHLAL HEDKPDFVGI ICT