

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBC448Mu01 100µq

Recombinant Dystrophin Associated Glycoprotein 1 (DAG1)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: His28~Pro406

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: Q62165

Host: E. coli **Purity: >90%**

Endotoxin Level: <1.0EU per 1µg (determined by

the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose, 0.01% sarcosyl.

Predicted isoelectric point: 6.4

Predicted Molecular Mass: 44.1kDa

Accurate Molecular Mass: 49kDa as determined by

SDS-

The possible reasons that the actual band size differs from the predicted are as follows:

PAGE reducing conditions.

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.) Note:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.



- TEL: 4006-871-227 Web: www. ybio. net Email: shybio@126. com 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

HWP SEPSEAVRDW KNQ LEASMHS VLSDFQEAVP TVVGIPDG TA VVG RSFRVSI PTDLIASSGE IIKVSAAGKE ALPSWLHWDP HSHILEGLPL DTDKGVHYIS VSAARLGANG SHVPQTSSVF SIEVYPEDHN EPQSVRAASS DPGEVVPSAC AADEPVTVLT VILDADLTKM TPKQRIDLLN RMQSFSEVEL HNMKLVPVVN NRLFDMSAFM AGPGNAKKVV ENGALLSWKL GCSLNQNSVP DIRGVETPAR EGAMSAQLGY PVVGWHIANK KPTLPKRLRR QIHATPTPVT AIGPPTTAIQ EPPSRIVPTP TSPAIAPPTE TMAPPVRDPV PGKPTVTIRT RGAIIQTPTL GPIOPTRVSE AGTTVPGOIR PTLTIP