YBB983Ra01 50µg

Recombinant Microtubule Associated Protein Tau (MAPt)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

> 9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Ala34 Ser368 (Accession # P19332), with two N-terminal Tags, His-tag and T7-tag.

Host: E. coli

Subcellular Location: Cytoplasm, cytosol. Cell

membrane; Peripheral membrane protein; Cytoplasmic

side. Cytoskeleton. Cell projection, axon.

Purity: >95%


Endotoxin Level: <1.0EU per 1µg (determined by the

LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris 500mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01%

sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point:



The possible reasons that the actual band size differs from the predicted are as follows: 9.9 Predicted Molecular Mass:

38. 5kDa

Accurate Molecular Mass: 48kDa as determined by SDS-PAGE reducing conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in ddH2O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with two N-terminal Tags, His-tag and T7-tag, its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- AEEAGIG DTPNMEDQAA GHVTQARVAG VSKDRTGNDE KKAKGADGKT GAKIATPRGA ATPGQKGTSN ATRIPAKTTP



SPKTPPGSGE PPKSGERSGY SSPGSPGTPG SRSRTPSLPT PPTREPKKVA VVRTPPKSPS ASKSRLQTAP VPMPDLKNVR SKIGSTENLK HQPGGGKVQI INKKLDLSNV QSKCGSKDNI KHVPGGGSVH IVYKPVDLSK VTSKCGSLGN IHHKPGGGQV EVKSEKLDFK DRVQSKIGSL DNITHVPGGG NKKIETHKLT FRENAKAKTD HGAEIVYKSP VVSGDTSPRH LSNVSSTGSI DMVDSPQLAT LADEVSAS

[REFERENCES]

- 1. Goedert M., et al. (1992) Proc. Natl. Acad. Sci. U.S.A. 89:1983-1987.
- 2. Georgieff I.S., et al. (1993) J. Cell Sci. 105:729-737.
- 3. Sadot E., et al. (1994) J. Mol. Biol. 241:325-331.
- 4. Kosik K.S., et al. (1989) Neuron 2:1389-1397.