



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YB91461Hu01 50 $\mu$ g

Azurocidin 1 (AZU1)

Organism: Homo sapiens (Human)

*Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

7th Edition (Revised in May, 2013)

## [ PROPERTIES ]

**Residues:** Pro23~Gly247 (Accession # P20160), with N-terminal His-Tag.

**Host:** *E. coli*

**Subcellular Location:** Cytoplasmic granule.

**Purity:** >95%

**Endotoxin Level:** <1.0EU per 1 $\mu$ g (determined by the LAL method).

**Formulation:** Supplied as lyophilized form in PBS, pH7.4, containing 5% sucrose, 0.01% sarcosyl.

**Predicted isoelectric point:** 9.1

**Predicted Molecular Mass:** 25.9kDa


**Accurate Molecular Mass:** 30kDa as determined by SDS-PAGE reducing conditions.

**Applications:** SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

**Note:** The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

## [ USAGE ]

Reconstitute in sterile PBS, pH7.2–pH7.4.

## [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2–8°C for one month.

Aliquot and store at –80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.




## [ SEQUENCES ]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MG H H H H H S G S E F - P L L D I V G G R K A R P R Q F P F L A S I Q N Q G R H F C G G A L I H A R  
FVMTAASCFQ SQNPGVSTVV LGAYDLRRRE RQSRQTFSIS SMSENGYDPQ QNLNDMLLQ  
LDREANLTSS VTILPLPLQN ATVEAGTRCQ VAGWGSQRSG GRLSRFPRFV NVTVTPEDQC  
RPNNVCTGVL TRRGICNGD GGTPLVCEGL AHGVASFSLG PCGRGPDDFT RVALFRDWID  
GVLNPG

## [ REFERENCES ]

1. Daigo K., *et al.* (2012) Mol. Cell Proteomics 11:M111.015073–M111.015073.
2. Chew M.S., *et al.* (2012) Inflamm. Res. 61:375–379.
3. Schou M., *et al.* (2011) Innate Immun 17:60–69.
4. Linder A., *et al.* (2010) J. Invest. Dermatol. 130:1365–1372.

**Unique product Superb quality Client favorite Nicest service**  ISO9001:2008;  ISO13485:2003; 

11271 Richmond Avenue, H104, Houston, TX 77082, USA | Toll free: 001-888-960-7402 | Fax: 001-832-538-0088 | Email: mail@uscnk.us | Http://www.uscnk.us  
Export Processing Zone, Wuhan 430056, P.R.China | Toll free: 0086-800-880-0687 | Fax: 0086-27-8425-9551 | Email: mail@uscnk.com | Http://www.uscnk.com