



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB329Hu01 10 μ g

Recombinant Microtubule Associated Protein 2 (MAP2)

Organism Species: Homo sapiens (Human)

*Instruction
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan,
2014)

[PROPERTIES]

Residues: Leu727~Asp985

Tags: N-terminal His-Tag

Accession: P11137

Host: *E. coli*

Subcellular Location: Cytoplasm, cytoskeleton.

Purity: >95%

Endotoxin Level: <1.0EU per 1 μ g (determined by
the LAL method).

Formulation: Supplied as lyophilized form in PBS,
pH7.4, containing 5% sucrose, 0.01% sarcosyl.


Predicted isoelectric point:

The possible reasons that the actual band size differs from the predicted are as follows:

4.7 Predicted Molecular Mass:

29.7kDa

Accurate Molecular Mass: 43kDa as determined by SDS-PAGE reducing
conditions. Applications: SDS-PAGE; WB; ELISA; IP.





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

(May be suitable for use in other assays to be determined by the end user.)

Note:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in sterile ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

LAS D ILT N TS G SMD EG D D YL PATT PALE KA P C FP VE SK E EE Q IE KV KAT G EE S T
QAEISCESPF LAKDFYKNGT VMAPDLPEML DLAGTRSRLA SVSADAEVAR RKSVPSETVV
EDSRTGLPPV TDENHVIVKT DSQLEDLGYC VFNKYTVPLP SPVQDSENLS GESGTFYEGT



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

DDKVRRLAT DLSLIEVKLA AAGRVKDEFS VDKEASAHIS GDKSGLSKEF DQEKKANDRL
DTVLEKSEEH ADSKEHAKKT EEAGD

[RELEVANCE]

1. Albala J. S., *et al.* (1993) *Gene* 136:377-378.
2. Dammerman M., *et al.* (1989) *J. Neurosci. Res.* 24:487-495.
3. Kosik K. S., *et al.* (1988) *J. Neurochem.* 51:587-598.
4. Zamora-Leon S. P., *et al.* (2005) *J. Biol. Chem.* 280:1962-1970.