

TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA547Si01 100µg

Recombinant Vascular Cell Adhesion Molecule 1 (VCAM1)

Organism Species: Rhesus monkey (Simian)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Leu1~Asp81

Tags: Two N-terminal Tags, His-tag and T7-tag

Accession: Q58T13

Host: E. coli

Purity: >90%

Endotoxin Level: <1.0EU per 1 µ g

(determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM

Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM

DTT, 0.01% sarcosyl, 5% trehalose, and preservative.

Predicted isoelectric point: 5.5


Predicted Molecular Mass: 13.2kDa

Applications: SDS-PAGE: WB: ELISA: IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile ddH2O.





[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below. LDRLEIELLK GETVLENKEF LEDTDMKSLE NKSLEATFIP TTEDTGKALV CQAKLHMDEM EFEPKQRQST QTLYVNVAPR D

[REFERENCES]

- 1. Seachord C.L., et al. (2005) Biol. Reprod. 72:1305-1314.
- 2. Cybulsky M., et al. (1991) Cytogenet. Cell Genet. 58: 1852.
- 3. Barreiro O., et al. (2002) J. Cell Biol. 157 (7): 1233 45.
- 4. Eibl RH., Benoit M. (2004) IEE Proc Nanobiotechnol. 151 (3): 128-32.