TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBA007Bo01 10µg

Recombinant Angiogenin (ANG)

Organism Species: Bos taurus; Bovine (Cattle)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

11th Edition (Revised in May, 2016)

[<u>PROPERTIES</u>]

Source: Prokaryotic expression. Host: *E. coli*

Residues: Ala24~His148

Tags: N-terminal His-Tag

Tissue Specificity: Liver.

Subcellular Location: Secreted.

Purity: >98%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA,

1mM DTT, 0.01% sarcosyl, 5%Trehalose and Proclin300.

Original Concentration: 200 ug/mL

Applications: SDS-PAGE; WB; ELISA; IP; CoIP; ReporterAssays;

Purification; Amine Reactive Labeling.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.2

Predicted Molecular Mass: 15.9kDa

Accurate Molecular Mass: 18kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80° C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at $37^\circ C$ for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCE</u>]

AODDYRY IHFLTOHYDA KPKGRNDEYC FNMMKNRRLT RPCKDRNTFI HGNKNDIKAI CEDRNGOPYR GDLRISKSEF OITICKHKGG SSRPPCRYGA TEDSRVIVVG CENGLPVHFD ESFITPRH




Figure 1. Gene Sequencing (Extract)






Figure 2. SDS-PAGE

Bio^{在增生物}T

EL:4006-871-227 Web:www.ybio. net Email:shybio @126.com