



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBF219Mu01 10 $\mu$ g

Recombinant Ectonucleotide Pyrophosphatase/Phosphodiesterase 1 (ENPP1)

Organism Species: *Mus musculus* (Mouse)

*Instruction  
manual*

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

## [ PROPERTIES ]

Residues: Gly568~Asp789 (Accession # P06802), with two N-terminal Tags, His-tag and GST-tag.

Host: *E. coli*


Subcellular Location: Cell membrane; Secreted.

Single-pass type II membrane protein. Basolateral cell membrane.

Purity: >95%

Endotoxin Level: <1.0EU per 1 $\mu$ g (determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris, 500mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative.





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Predicted isoelectric point:

6.4 Predicted Molecular Mass:

58.3kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

### [ USAGE ]

Reconstitute in ddH<sub>2</sub>O.

### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

### [ SEQUENCES ]

The target protein is fused with two N-terminal Tags, His-tag and GST-tag, its sequence is listed below.

MSPILGYWKI KGLVQPTRLL LEYLEEKYEE HLYERDEGDK WRNKKFELGL EFPNLPYYID  
GDVKLTQSM A IIRYIADKHN MLGGCPKERA EISMLEGAVL DIRYGVSR IA YSKDFETLKV  
DFLSKLP EML KMFEDRLCHK TYLNGDHVTH PDFMLYDALD VVLYMDPMCL DAFPKLVCFK  
KRIEAIPQID KYLKSSKYIA WPLQGQWATF GGGDHPPKSD GSTSGSGHHH HHHSAGLVPR  
GSTAIGMKET AAKAFERQHM DSPDLGTGGG SGIEGRGSMG YRGSEF-GSH GSLNHLKKP  
IYNPSHPKEE GFLSQCP IKS TSNDLGCTCD PWIVPIKDFE KQLNLTEDV DDIYHMTVPY



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

GRPRILLKQH HVCLLQQQF LTGYSLDLLM PLWASYTFLR NDQFSRDDFS NCLYQDLRIP  
LSPVHKCSYY KSNKLSYGF LTPPRLNRVS NHIYSEALLT SNIVPMYQSF QVIWHYLHDT  
LLQRYAHERN GINNVSGPVF DFDYDGRYD

## [ REFERENCES ]

1. van Driel I.R., Goding J.W. (1987) J. Biol. Chem. 262:4882-4887.
2. Rebbe N.F., *et al.* (1991) Proc. Natl. Acad. Sci. U.S.A. 88:5192-5196.
3. Banakh I., *et al.* (2002) Eur. J. Immunogenet. 29:307-313.
4. Belli S.I., *et al.* (1993) Eur. J. Biochem. 217:421-428.