TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

YBB252Ra01 10µg

Androgen Receptor (AR)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

kDa 66.2

45

33

26

20

14.4

# [ PROPERTIES ]

Residues: Val491<sup>Ser679</sup> (Accession # P15207), with Nterminal His-Tag. Host: E. coli Subcellular Location: Nucleus. Cytoplasm. Purity: >95% Endotoxin Level:  $\langle 1.0EU \text{ per } 1 \mu g \rangle$  (determined by the LAL method). Formulation: Supplied as lyophilized form in PBS, pH7.4, 15% SDS-PAGE containing 5% sucrose, 0.01% sarcosyl. Predicted isoelectric point:

The possible reasons that the actual band size differs from the predicted are as follows: 7.1 Predicted Molecular Mass: 22. 5kDa

Accurate Molecular Mass: 25kDa as determined by SDS-PAGE reducing conditions. Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

<sup>9</sup>th Edition (Revised in Jul, 2013)



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Note:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

## [ USAGE ]

Reconstitute in sterile PBS, pH7.2-pH7.4.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

### SEQUENCES ]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSGSEF-VNRVPYPSPS CVKSEMGPWM ENYSGPYGDM RLDSTRDHVL PIDYYFPPQK TCLICGDEAS GCHYGALTCG SCKVFFKRAA EGKQKYLCAS RNDCTIDKFR RKNCPSCRLR KCYEAGMTLG ARKLKKLGNL KLQEEGENSS AGSPTEDPSQ KMTVSHIEGY ECQPIFLNVL EAIEPGVVCA GHDNNQPDS



TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

## [ <u>REFERENCES</u> ]

- 1. Stanworth R.D., et al. (2008) Eur. J. Endocrinol. 159:739-746.
- 2. Li R. J., et al. (2008) Apoptosis 13:959-971.
- 3. Taylor A.H., et al. (2008) Br. J. Dermatol. 158:319-328.
- 4. Albertelli M.A., et al. (2008) Hum. Mol. Genet. 17:98-110.