YBC117Mu01 100μg

Recombinant Activin A Receptor Type II A (ACVR2A)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Ala20~Lys95 linked with TALKCTFVAV

RAICVMKSPL IFRRWKSHSP LQILLHRSHP

DSSTTTTTTE IRLLTKPERK LSWLLPPLSN N Tags: N-

terminal His-Tag

Accession: P27038

Host: E. coli

Subcellular Location: Membrane; Single-pass

type I membrane protein.

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL

method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% trehalose, 0.01% sarcosyl.


Predicted isoelectric point: 5.6

Predicted Molecular Mass: 15.0kDa

Accurate Molecular Mass: 19kDa as determined by SDS-PAGE reducing conditions.

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)





TEL:4006-871-227 Web:www.ybio.net Email:shybio@126.com

Note: The possible reasons that the actual band size differs from the predicted are as follows:



- TEL: 4006-871-227 Web: www.ybio.net Email: shybio@126.com 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

A IL G R S E T Q E C LF F N A N W E R D RT N Q T G V E P C YG D K D K R R H C FAT W K N I S G S IEIVK QG CWL DDIN CY DRTD CIEK K TALK CTFVAV RAIC VMKSPL IFRRWK SHS P LOILLHRSHP DSSTTTTTTE IRLLTKPERK LSWLLPPLSN N