

YBA049Mu01 50μg Recombinant Interferon Gamma (IFNg) Organism Species: Mus musculus (Mouse) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: His23~Cys155

Tags: N-terminal His-Tag

Accession: P01580

Host: E. coli

Subcellular Location: Secreted.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL

method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 1mM DTT, 5% trehalose, and

preservative.

Predicted isoelectric point: 8.8


Predicted Molecular Mass: 17.0kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.





[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

HGTVIESL ESLNNYFNSS GIDVEEKSLF LDIWRNWQKD GDMKILQSQI ISFYLRLFEV LKDNQAISNN ISVIESHLIT TFFSNSKAKK DAFMSIAKFE VNNPQVQRQA FNELIRVVHQ LLPESSLRKR KRSRC

[REFERENCES]

- 1. Goshima N., et al. (2008) Nat. Methods 5:1011-7.
- 2. The MGC Project Team (2004) Genome Res. 14:2121-2127.
- 3. Rinderknecht E., et al. (1984) J. Biol. Chem. 259:6790-7.
- 4. Gray P.W., Goeddel D.V. (1983) Proc. Natl. Acad. Sci. U.S.A. 80:5842-5846.
- 5. Landar A., et al. (2000) J. Mol. Biol. 299:169-79.
- 6. Carninci P., et al. (2005) Science 309:1559-1563.
- 7. Dufour C., et al. (2004) Br. J. Haematol. 126:682-5.